

ISSN 2509-4327 (print)
ISSN 2510-4780 (online)

Inter
GING



Deutscher Wissenschaftsherold German Science Herald

№ 4/2017

Die Zeitschrift „Deutscher Wissenschaftsherold“ ist eine Veröffentlichung mit dem Ziel ein breites Spektrum der Wissenschaft allgemeinverständlich darzustellen. Die Redaktionsleitung versteht sich als Vermittler zwischen Wissenschaftlern und Lesern. Durch die populärwissenschaftliche Bearbeitung wird es möglich unseren Lesern neue wissenschaftliche Leistungen am besten und vollständigsten zu vermitteln. Es werden Untersuchungen, Analysen, Vorlesungen, kurze Berichte und aktuelle Fragen der modernen Wissenschaft veröffentlicht.

Impressum

Deutscher Wissenschaftsherold – German Science Herald

Wissenschaftliche Zeitschrift

Herausgeber:

InterGING

Sonnenbrink 20

31789 Hameln, Germany

Inhaber: Marina Kisiliuk

Tel.: + 49 51519191533

Fax.: + 49 5151 919 2560

Email: info@dwherold.de

Internet: www.dwherold.de

Chefredakteur/Editor-in-chief:

Marina Kisiliuk

Korrektur:

O. Champela

Gestaltung:

N. Gavrilets

Auflage: № 4/2017 (September) – 30

Redaktionsschluss September, 2017

Erscheint vierteljährlich

Editorial office: InterGING

Sonnenbrink 20

31789 Hameln, Germany

Tel.: + 49 51519191533

Fax.: + 49 5151 919 2560

Email: info@dwherold.de

Deutscher Wissenschaftsherold - German Science

Herald is an international, German/English language,

peer-reviewed, quarterly published journal.

№ 4/2017

Passed in press in September 2017

Druck: WIRMachenDRUCK GmbH

Mühlbachstr. 7

71522 Backnang

Deutschland

Der Abdruck, auch auszugsweise, ist nur mit ausdrücklicher Genehmigung der InterGING gestattet. Die Meinung der Redaktion oder des Herausgebers kann mit der Meinung der Autoren nicht übereinstimmen. Verantwortung für die Inhalte übernehmen die Autoren des jeweiligen Artikels.

INDEXING: Google Scholar, WorldCat, Index Copernicus, InfoBase Index, Journal Index, Citefactor, International Scientific Indexing, JIFACTOR, Scientific Indexing Services, International Institute of Organized Research.



JIFACTOR



CiteFactor
Academic Scientific Journals



Scientific Indexing Services



INTERNATIONAL
Scientific Indexing



MIAR

http://miar.ub.edu/issn/2509-4327

© InterGING

© Deutscher Wissenschaftsherold – German Science Herald

REDAKTIONSKOLLEGIUM / INTERNATIONAL EDITORIAL BOARD:

Jurga Bernatoniene, Dr., Prof.
Physics Lithuania
jurgabernatoniene@yahoo.com

Arvidas Galdikas, Dr. habil., professor
Physics Lithuania,
arvidas.galdikas@ktu.lt

Kristina Ramanauskienė, Ph.dr., Prof.
Pharmacy, Lithuania
kristinaraman@gmail.com

Khpaliuk Alexander, Dr. med. habil., Prof.
Pharmakologie, Belarus
clinicfarm@bsmu.by

Arnold M. Gegechkori, Dr., full Prof.
Biology, Georgia
arngegechkori@yahoo.com

Omari Mukbaniani, Prof., DSc.
Chemistry, Georgia
omar.mukbaniani@tsu.ge

Teimuraz Lezhava, Prof.
Genetics, Georgia
teimuraz.lezhava@tsu.ge

Shota A. Samsoniya, Prof.
Chemistry, Georgia
shota.samsonia@tsu.ge

Mdzinarashvili Tamaz, DSc., Prof.
Biophysics, Georgia
tamaz.mdzinarashvili@tsu.ge

Aliaksandr V.Prokharau, MD, PhD, MSc Prof.
Oncology, Belarus
aprokharau@gmail.com

Pyrochkin V., MD, PhD, MSc Prof.
Theraphy, Belarus
wlad_cor@mail.ru

Golubev A.P., BD, Prof.
Ecology, Belarus
algiv@rambler.ru

Makarevich A., MD, PhD, Prof.
Theraphy, Belarus
makae@bsmu.by

Kanunnincova N., BD, Prof.
Physiology, Belarus
n.kanunnikova@grsu.by

Giedrius Vanagas, Prof.
Internal Medicine, Lithuania
Giedrius.Vanagas@lsmuni.lt

Armuntas Baginskas, Prof.
Neurofiziologija, Lithuania
Armuntas.Baginskas@lsmuni.lt

Ricardas Radisauskas, MD., Ph.D., Prof.
Cardiology, Lithuania
Ricardas.Radisauskas@lsmuni.lt

Meyramov Gabit, Prof.
Cytology and Histology, Kazakhstan
meyramow@mail.ru

Aisha Mohammed Abd al-salam Shahlol
Ph.D. in Medical Bacteriology, Libya
Ais.shahlol@sebhau.edu.ly

Edmundas Kadusevicius, MD, PharmD, PhD, Prof.
Pharmacology, Lithuania
Edmundas.Kadusevicius@lsmuni.lt

Ivo Grabchev, Prof., PhD.
Chemistry, Bulgaria
i.grabchev@chem.uni-sofia.bg
grabchev@mail.bg

Mariyana Ivanova Lyubenova, Prof., PhD.
Ecology, Bulgaria
ryann@abv.bg
ryana_1@yahoo.com

Tsvetanka Tsankova Marinova, MD, PhD, DMedSci,
Biologv. Bulgaria
tmarinova@yahoo.com

Evgueni D. Ananiev, Prof PhD,
Biology. Bulgaria
evgueni_ananiev@yahoo.com

Plamen G. Mitov, Prof., PhD.
Biology, Bulgaria
mitovplamen@gmail.com

Atanas Dimov Arnaudov, Ph.D.
Physiology, Bulgaria
arny87@yahoo.co.uk

Iliana Georgieva Velcheva, PhD,
Ecology, Bulgaria
anivel@abv.bg

Osman Demirhan, Prof.
Biology, Turkey
osdemir@cu.edu.tr

Jharna Ray, M. Sc., PhD, Prof.
Neurogenetics, India
Indijharnaray@gmail.com

Marián Halás doc. RNDr, Ph.D.
Human geography, Czech
marian.halas@upol.cz

Ayfer Pazarbasi Prof.Dr.
Biology, Turkey
payfer@cu.edu.tr

Tusharkanti Ghosh Prof.
Physiology, India
tusharkantighosh53@yahoo.in

Khudaverdi Gambarov Gambarov, Prof.
Microbiology, Azerbaijan
khuda1949@mail.ru

Rovshan Ibrahimkhalil Khalilov, Prof.
Biophysics, Azerbaijan
hrovshan@hotmail.com

Svitlana Antonyuk, Dr.phil.
Stony Brook University, USA
Linguistics

Samuel M.Johnson, Prof.Dr.phil.
Theology, Wells, Maine, USA
djtjohnson@earthlink.net

Satanovsky Leon MD/PhD.
Perio-odontologie, Israel
satleonid@gmail.com

Lists of references are given according to the Vancouver style

**Pecheryaga S.V.,
Marinchina I.M.**

Department of Obstetrics, Gynecology and Perinatology, Higher State Educational Establishment of Ukraine "Bukovinian State Medical University", Chernivtsi, Ukraine

FEATURES OF HEMODYNAMIC CHANGES IN SPIRAL ARTERIES WITH LOW PLACENTATION AT THE EARLY GESTATIONAL AGE

Abstract. *The article presents the results of Doppler studies in spiral arteries of indices of resistance and blood flow velocities in the I trimester of gestation in pregnant women with low placentation. As a result of the studies, a significant increase in resistance and a decrease in blood flow rates in early pregnancy in pregnant women with low placentation was found in comparison with normal placement of the chorion, which may be a factor of high risk of primary placental dysfunction development in the future.*

Key words: *low placentation, I trimester of gestation, Doppler examination, spiral arteries.*

Introduction. Among the causes that negatively affect the course of pregnancy and the state of the fetoplacental system, the anomalies of the implantation of the fetal egg play an important role, in particular, low placentation accompanied by insufficient functioning of the uterine-placental complex, which is caused by the character of the vascularization of the lower parts of the uterus and a decrease in the placental blood flow [2, 3]. The adequate functioning of the mother-placenta-fetus system depends primarily on the process of implantation of the fetal egg, cytotrophoblastic invasion, as well as the subsequent transformation of the spiral arteries. The results of studies carried out in recent years have shown that the formation of intrauterine suffering is laid precisely in the early gestation period, when problems in the body of a woman, the condition of the endo- and myometrium determines the inferior formation of the embryo, fetus and extraembryonic formations: amniotic fluid, umbilical cord, placenta, placental bed. Disruptions in the formation of the vascular system of the mucous membrane of the uterus is accompanied by structural changes of the latter; conditions for implantation of the fetal egg are significantly reduced. In this regard, a noninvasive study of the endometrium during blastocyst implantation is important [5]. The method of Doppler investigation of blood flow in the fetoplacental system is safe, relatively simple and at the same time highly informative in assessing its functional reserves [1, 4].

The leading role in the pathogenesis of placental dysfunction is played by disorders of uterine-placental circulation.

The constancy of the uterine-placental blood flow ensures a gestational reorganization of the spiral arteries. The physiological changes experienced by spiral arteries with the progression of uncomplicated pregnancy are characterized by elastolysis, degeneration of the muscular layer and replacement of the muscular and elastic fibrinoid shell with the expansion of the artery lumen [3].

Objective of the study. Conduct an evaluation of Doppler blood flow in the spiral arteries with low placentation.

Materials and methods. The main group of the study included pregnant women in the period from 5 to 12 weeks of pregnancy with a low location of the chorion (50 pregnant women). The control group consisted of 50 pregnant women at the same gestational age with the arrangement of the chorion in the uterine body and fundus. All the examined women of the primary and control groups were divided into 2 subgroups, depending on the gestation period (25 pregnancies in the period of 5-8 weeks of gestation and 25 women in the period of 9-12 weeks of gestation). The groups studied were representative of age, height and weight, and social employment. All were subjected to ultrasound on the apparatus «Voluson Exspert 730», to all women the location of the chorion and the fetal egg, the crown-rump length of the fetus were determined, as well as

the Doppler examination of the blood flow in the spiral arteries of the uterus. Indices of vascular resistance were calculated: the systolic-diastolic ratio (SDR), the pulsation index (PI), the resistance index (RI), as well as the pulse systolic rate (PSR), the end diastolic velocity (DV), the average distal velocity (ADV), the peak speed averaged over time (PSAT). Statistical processing of the obtained indicators was carried out by determining the Student's criteria.

Results of the study and their discussion.

When analyzing the data obtained, it was found that in pregnant women with a low chorion placement at 5-8 weeks of gestation with Doppler study in spiral arteries, all resistance indices were higher: SDR- 3.9 ± 0.3 , PI- 0.71 ± 0.04 , RI - 1.4 ± 0.1 compared to the control: SDR- 3.0 ± 0.3 , PI- 0.58 ± 0.03 , RI- 1.1 ± 0.08 (p0.05). Accordingly, blood flow velocities were lower in the main group, but the only significant difference was in the PSAT index, which in the main group was 19.5 ± 1.2 cm / s, and in the control group - 29.6 ± 1.8 cm / s.

This may indicate an imperfect transformation of the walls of the spiral arteries in the abnormal position of the chorion, disorders of the first wave of invasion of the cytotrophoblast and the disrupted blood supply of the emerging placental bed. In turn, this determines the limit of utero-placental perfusion, the inadequacy of blood flow in the intervillaceous space, where the blood flows from the spiral arteries, the stasis of blood in it and the disruption of the gas exchange process.

When evaluating Doppler indices in spiral arteries in 9-12 weeks of pregnancy, high peripheral resistance values remain in the main group, respectively, SDR- $3,1 \pm 0,4$; PI 0.64 ± 0.05 ; RI- $1,1 \pm 0,11$ in comparison with the control of $2,1 \pm 0,2$; 0.5 ± 0.04 ; 0.82 ± 0.04 , respectively (p0.05). SDR rates by 32.2%, PI by 21.9%, and RI by 25.5% higher in women with low placentation compared with pregnant women with normal placement of chorion. So, we see that there is a preservation of high preplacental resistance of the blood flow after passing the "peak" of the first wave of involution of the cytotrophoblast.

The decrease in blood flow rates is more significant in the spiral arteries during 9-12 weeks of gestation, which is manifested by a significant decrease in SDR- $25,4 \pm 1,4$ cm / s, ADV- $24,2 \pm 1,4$

cm / s, PSAT- $38,1 \pm 1.6$ cm / sec in pregnant women with low chorion distribution compared with control group, where the corresponding indices make up 33.7 ± 1.5 cm / s, 34.4 ± 1.4 cm / s, $56.3 \pm 2, 1$ cm / sec (p0.05).

Consequently, the disruption of blood flow in spiral arteries with low placentation progresses in the dynamics of pregnancy, which leads to a deepening of placental dysfunction and an increase in the level of complications of pregnancy.

Conclusions. 1. With low placentation in the I trimester of gestation the formation of highly resistant blood flow and the decrease in blood flow rates occur in the spiral arteries.

2. The revealed changes in hemodynamics in spiral arteries with low placement of chorion may be the factors of high risk of development of primary placental dysfunction in the future.

Prospects for further research. To study the parameters of volume blood flow in the chorion with abnormal placentation.

References:

1. Markin LB, Shatilovich KL, Nadorshina NE. *Doplerometrija v akusherstvi: gemodinamichni osoblivosti funkcional'noi sistemi mati-placenta-plid. Reproaktivnoe zdorov'e zhenshhiny. 2007; 30(1):36-9.*
2. Minkina JeR. *Nizkaja placencija kak faktor riska pri beremennosti. Vestnik Rossijskogo gosudarstvennogo medicinskogo universiteta. 2008; (3):104.*
3. Ponomareva NA. *Prognosticheskie kriterii razvitija placentarnoj nedostatochnosti v I trimestre beremennosti. Mater 7-go Ross Forum Mat' i ditja. M; 2005. P. 208-9.*
4. Titchenko LI, Chechneva MA, Zhukova NV. *Trehmernaja rekonstrukcija v I trimestre beremennosti. Rossijskij vestnik akushera-ginekologa. 2003; (5):16-20.*
5. Kofinas AG, Pentry M, Greiss FC. *The effect of placental location on uterine flow velocity waveforms. Amer J Obster Gincol. 2010; 159(6):1504-8.*
7. Brushlinskogo AV, Shapiro AZ, editors. *Pezeshkian N. Psihosomatika i pozitivnaja psihoterapija. M: Institut pozitivnoj psihoterapii; 2006. 464 p.*

CONTENT:

Galagdina A.A., Dmytrenko R.R., Bambuliak A.V. Diagnostics of ischemic-reperfusion damage of the brain in rats afflicted with diabetes mellitus	3
Guranych S.P., Voronych-Semchenko N.M., Guranych T.V. Macro- and microelement status of rats with insulin resistance against the ground of iodine deficiency	6
Fedyshyn T.V., Maliar V.V., Maliar V.A. Peculiarities of utero-placental blood circulation formation in women with spontaneous and recurrent miscarriages associated with vagina dysbiosis	10
Rusnak V.F., Bedyk V.V. Growth of the pharynx at the end of the fetal stage of human ontogenesis	13
Teplytskyi S.S. Formation and development of the skin on the palmar surface of the hand throughout the period of prenatal ontogenesis and its importance in dermatoglyphics	16
Tkachuk N.P., Bilookyi V.V., Gyrla Ya.V., Sheremet M.I. Evaluating the efficiency of the scale for prediction of post-operational relapse in patients with nodular goiters	20
Yemelyanenko N.R. Anatomical transformations of the nasal septum in childhood	24
Kavun M.P. Morphogenesis of the hepatic-duodenum ligament in early ontogenesis of the human	26
Kotyuzhinskaya S.G., Umansky D.A. Functional state of lipittransport system in patients with atherosclerosis with fatty load	28
Lomakina Yu.V., Burdeina M.P. Stress-associated changes in the excretory function of the kidneys in old rats under the conditions of a usual light period	32
Malyar V.V. Structural and functional features of fetal membranes in pregnant women with moderate idiopathic oligo- and polyhydramnios	35
Nesterak R.V., Gasyuk M.B. Pilot investigation of the method of interactive training of patients at the stage of medical rehabilitation and treatment	38
Pecheryaga S.V., Marinchina I.M. Features of hemodynamic changes in spiral arteries with low placentation at the early gestational age	42
Psychenko V.V., Chernov V.S., Frenkel Yu.D. The status of extraorganic blood flow in pineal gland of rats under conditions of acute stress and twenty-four hour darkness	44
Reshetilova N.B., Glubochenko O.V., Kulish N.M., Dudko A.G. Formation of anterior cerebral vesicle cavities at the 5th week of the embryonic period	47
Riznichuk M.O., Galitskaya V.O., Dyhodyuk Yu.V., Kravchuk Yu.V., Vakaryuk O.V. Prader-will syndrome, diagnostics and currency features	50
Shalamay U.P., Pavlikivska B.M., Voronich-Semchenko N.M. The state of autonomous heart regulation in adolescents with light iodine deficiency and latent iron deficiency	52
Shutova N.A., Nikolayeva O.V., Sulkhodost I.O. Electromagnetic radiation impact on the cellular protective mechanisms in experiment	58
Yasnikovska S.M., Hrytsak H. Evaluation of clinic-laboratory and ultrasonic research results in different forms of the chorion's pathology in the first three-month of gestation	61
Yashchyshyn Z.M., Zaiats L.M., Yurkiv I.Y., Maslyak K.T., Vodoslavskaya N.Y., Sikomas M.T. Changes in neuroglial interrelation of muscle-intestinal nerve plexus of esophagus after one-sided crossing of vagosympathetic trunk	64
Navarchuk N.M., Kosteniuk S.V. Morphogenesis of the dentognathic apparatus during the early times of the human ontogenesis	67
Rusnak V.F., Bedyk V.V. Features of pharyngeal morphogenesis in five-week embryos	70
Talanova O.S., Apt O.A. Specifics of distribution of glycosaminoglycans in the white pulp of the spleen and stroma of rats after experimental modeling injection inside the fetus of antigens of different nature	72
Pivtorak K.V., Mazur I.A., Voloshin M.A. Correction of metabolic disorders caused by non-alcoholic fatty liver disease	74
Rozhko V.I. Research of content correlation of immunoglobulins and lisozyme in oral fluid of children with rampant caries against the background of gastro-intestinal diseases	78
Karavan Ya.R., Havaleshko V.P. Up-to-date anesthetic possibilities in dentistry practice in diagnosis of the patient's allergic status	80



Deutscher Wissenschaftsherold German Science Herald

**Bibliographic information published by the Deutsche Nationalbibliothek
The Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie; detailed
bibliographic data are available on the Internet at <http://dnb.dnb.de>**

**№ 4/2017 – 30
Passed in press in September 2017**



WirmachenDruck.de

Sie sparen, wir drucken!